


PhenoBR Workshop


Probing the Top-Higgs sector with clustering techniques

Ricardo D'Elia Matheus




IFT - UNESP
INSTITUTO DE FÍSICA TEÓRICA

Obligatory “problems of the SM slide”



Are we done?

Obligatory “problems of the SM slide”



Dark matter?

Flavor Structure?

(most parameters of the SM)

Hierarchy of Scales?

Neutrino Masses?

(are they Dirac \rightarrow New DoF
are they Majorana \rightarrow New scale)

Baryogenesis?





Strong CP? Axions?

Gravitons?

Particle physics is only “over” if you are satisfied with: “I don’t know”

Beyond SM

The solution to most of these shortcomings involve new models:

- UV Complete, detailed models  Theoreticians used to love these
(e.g.: SUSY)
- Simplified models  Most experimental constraints apply here
- Effective models (inspired by some UV completion)
(e.g.: Composite Higgs Model)  We will talk about the Minimal Composite Higgs Model (MCHM soon)
- Model Independent EFT  All the rage nowadays (theoreticians and increasingly experimentalists too)

A common problem

Model parameter space is multidimensional

$$\mathcal{L} = \mathcal{L}_{\text{elem}} + \mathcal{L}_{\text{comp}}^{14} + \mathcal{L}_{\text{mix}}^{14} + \mathcal{L}_{\text{int}}^{14}$$


e.g.: MCHM₁₄

$$\mathcal{L}_{\text{int}}^{14} = -i \boxed{c_4} \bar{\Psi}_4 \not{D} \Psi_1 - i \boxed{c_9} \bar{\Psi}_9^{ij} \not{D}^i \Psi_4^j - i \frac{\boxed{c_{T9}}}{4\pi \boxed{f}} \bar{\Psi}_9^{ij} d_\mu^i d^{j\mu} \tilde{T} + \text{h.c.}$$

$$\begin{aligned} \mathcal{L}_{\text{mix}}^{14} &= f \text{Tr} \left[U^\top \bar{Q}_L^{14} U (\boxed{y_{L9}} \Psi_9 + \boxed{y_{L4}} \Psi_4 + \boxed{y_{L1}} \Psi_1) \right] + \text{h.c.} \\ &+ f \text{Tr} \left[U^\top \bar{T}_R^{14} U (\boxed{y_{R9}} \Psi_9 + \boxed{y_{R4}} \Psi_4 + \boxed{y_{R1}} \Psi_1) \right] + \text{h.c.} \end{aligned}$$

A common problem


Model parameter space is multidimensional



Model parameters

A common problem


Model parameter space is multidimensional...



... and not trivially connected to observables


A common problem

Model parameter space is multidimensional...



... and not trivially connected to observables


Common Solutions




It is the sound
approach!

... but we were not lucky

Common Solutions




Clustering



We want to group these points into **clusters** that have similar phenomenology

Clustering




What criteria to use?

Physical parameters


Mass Eigenstates

- Masses
- Widths
- Branching Ratios


Observables ✓



Clustering



Observables




We can also get a representative **benchmark point** for each cluster

These points can then be used to design search strategies or figure out constraints that apply across the cluster

Clustering Algorithm

As proposed in: A. Carvalho, M. Dall'Osso, T. Dorigo, F. Goertz, C. A. Gottardo and M. Tosi, JHEP 04 (2016) 126, arXiv: 1507.02245


We will do a bin by bin comparison of distributions obtained at different points in parameter space



Clustering Algorithm

As proposed in: A. Carvalho, M. Dall'Osso, T. Dorigo, F. Goertz, C. A. Gottardo and M. Tosi, JHEP 04 (2016) 126, arXiv: 1507.02245

We will do a bin by bin comparison of distributions obtained at different points in parameter space



P₁ vs. P₂


0 1000 3000 3000 4000 2000 6000

10⁰

Clustering Algorithm

As proposed in: A. Carvalho, M. Dall'Osso, T. Dorigo, F. Goertz, C. A. Gottardo and M. Tosi, JHEP 04 (2016) 126, arXiv: 1507.02245

We will do a bin by bin comparison of distributions obtained at different points in parameter space




SM vs. P_2

- SM
- MCHM₅, P_1
- MCHM₅, P_2
- MCHM₅, P_3

Clustering Algorithm

As proposed in: A. Carvalho, M. Dall'Osso, T. Dorigo, F. Goertz, C. A. Gottardo and M. Tosi, JHEP 04 (2016) 126, arXiv: 1507.02245

$$TS_{ab} = -2 \sum_{i=1}^{N_{bins}} \left[\log(n_{(i,a)!}) + \log(n_{(i,b)!}) - 2\log\left(\frac{n_{(i,a)} + n_{(i,b)}}{2}\right)! \right]$$



If $S_a = S_b \rightarrow TS_{ab} = 0$

$TS_{ab} < 0 \leftrightarrow S_a \neq S_b$
(increasingly so)



$TS_{ab} > TS_{cd}$ means
 S_a and S_b are more alike
than S_c and S_d

Clustering Algorithm

As proposed in: A. Carvalho, M. Dall'Osso, T. Dorigo, F. Goertz, C. A. Gottardo and M. Tosi, JHEP 04 (2016) 126, arXiv: 1507.02245

$$TS_{ab} = -2 \sum_{i=1}^{N_{bins}} \left[\log(n_{(i,a)}) + \log(n_{(i,b)}) - 2 \log \left(\frac{n_{(i,a)} + n_{(i,b)}}{2} \right) \right]$$


We can also sum over more than one kinematical distribution



Clustering Algorithm

As proposed in: A. Carvalho, M. Dall'Osso, T. Dorigo, F. Goertz, C. A. Gottardo and M. Tosi, JHEP 04 (2016) 126, arXiv: 1507.02245

Let the clustering begin!



Step 0


Simulate events for all points and get the kinematical distributions (build **samples**)

Each point will be its own cluster

Clustering Algorithm

As proposed in: A. Carvalho, M. Dall'Osso, T. Dorigo, F. Goertz, C. A. Gottardo and M. Tosi, JHEP 04 (2016) 126, arXiv: 1507.02245

Let the clustering begin!



Step 1

Compare all pairs of clusters


$$TS^{\min} = \min_{ab}(\{TS_{ab}\})$$

a and b run over all points in their cluster

Clustering Algorithm

As proposed in: A. Carvalho, M. Dall'Osso, T. Dorigo, F. Goertz, C. A. Gottardo and M. Tosi, JHEP 04 (2016) 126, arXiv: 1507.02245

Let the clustering begin!



Step 1

Compare all pairs of clusters

$$TS^{min} = \min_{ab}(\{TS_{ab}\})$$


a and b run over all points in their cluster

Merge the pair with the **highest** TS^{min}

Clustering Algorithm

As proposed in: A. Carvalho, M. Dall'Osso, T. Dorigo, F. Goertz, C. A. Gottardo and M. Tosi, JHEP 04 (2016) 126, arXiv: 1507.02245

Let the clustering begin!



Step 3

Compare all pairs of clusters

$$TS^{\min} = \min_{ab}(\{TS_{ab}\})$$


a and b run over all points in their cluster

Merge the pair with the **highest** TS^{\min}

Clustering Algorithm

As proposed in: A. Carvalho, M. Dall'Osso, T. Dorigo, F. Goertz, C. A. Gottardo and M. Tosi, JHEP 04 (2016) 126, arXiv: 1507.02245

Let the clustering begin!



Step n (final)

Criteria: avoid the extremes


N too big \rightarrow highly homogeneous clusters, unwieldy number

N too small \rightarrow highly heterogeneous clusters

Clustering Algorithm

As proposed in: A. Carvalho, M. Dall'Osso, T. Dorigo, F. Goertz, C. A. Gottardo and M. Tosi, JHEP 04 (2016) 126, arXiv: 1507.02245

Let the clustering begin!




Step n (final)

Criteria: in practice it is a bit subjective, we look for a step where decreasing N will merge two big unlike clusters

Clustering Algorithm

As proposed in: A. Carvalho, M. Dall'Osso, T. Dorigo, F. Goertz, C. A. Gottardo and M. Tosi, JHEP 04 (2016) 126, arXiv: 1507.02245

Benchmarks Points



For each of the clusters we can choose a point a maximizing:


$$TS_a^{\min} = \min_b(\{TS_{ab}\})$$

where b runs over all other points in that cluster

Clustering Algorithm

As proposed in: A. Carvalho, M. Dall'Osso, T. Dorigo, F. Goertz, C. A. Gottardo and M. Tosi, JHEP 04 (2016) 126, arXiv: 1507.02245

Benchmarks Points



For each of the clusters we can choose a point a maximizing:

$$TS_a^{min} = \min_b(\{TS_{ab}\})$$

where b runs over all other points in that cluster

Those are the benchmark points

Application to the Minimal Composite Higgs Model (MCHM)



C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)



pNGB of $SO(5) \rightarrow SO(4)$

New fermionic DoFs introduced in some representation of the $SO(5)$

MCHM₅: 5 of $SO(5) \rightarrow 4+1$ of $SO(4)$





Application to the Minimal Composite Higgs Model (MCHM)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

Mix with SM (3rd gen.): **Partial Compositeness**


MCHM₅: 5 of SO(5) → 4+1 of SO(4)



Application to the Minimal Composite Higgs Model (MCHM)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

Charge 2/3 sector



$$\mathcal{L}_{\text{comp}}^5 = \bar{\Psi}_4 i(\not{D} - i\phi)\Psi_4 - M_4 \bar{\Psi}_4 \Psi_4 + \bar{\Psi}_1 i\not{D}\Psi_1 - M_1 \bar{\Psi}_1 \Psi_1$$


$$\begin{aligned} \mathcal{L}_{\text{mix}}^5 = & f \bar{Q}_L^5 U [y_{L4}\Psi_4 + y_{L1}\Psi_1] + \text{h.c.} \\ & + f \bar{T}_R^5 U [y_{R4}\Psi_4 + y_{R1}\Psi_1] + \text{h.c.} \end{aligned}$$

$$\mathcal{M}_{2/3}^5 = \begin{bmatrix} 0 & \frac{1}{2}y_{L4}f(1 + \sqrt{1-\xi}) & \frac{1}{2}y_{L4}f(1 - \sqrt{1-\xi}) & \frac{1}{\sqrt{2}}y_{L1}f\sqrt{\xi} \\ -\frac{1}{\sqrt{2}}y_{R4}f\sqrt{\xi} & -M_4 & 0 & 0 \\ \frac{1}{\sqrt{2}}y_{R4}f\sqrt{\xi} & 0 & -M_4 & 0 \\ y_{R1}f\sqrt{1-\xi} & 0 & 0 & -M_1 \end{bmatrix}$$

Application to the Minimal Composite Higgs Model (MCHM)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

Charge 2/3 sector



Application to the Minimal Composite Higgs Model (MCHM)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

MCHM₅: $T^{(1)}$ $T^{(2)}$ $T^{(3)}$ B $X_{5/3}$

$f, |M_1|, |M_4|, \text{sign}(M_1), y_L$ and y_R

Fixed by the mass of the top

MCHM₁₄: 14 of SO(5) → 9+4+1 of SO(4)

7 top partners

$$M_{2/3}^{14} = \begin{bmatrix} 0 & \frac{1}{2}y_L f a_+ & -\frac{1}{2}y_L f a_- & -\frac{\sqrt{5}}{4}y_L f s_{2h} & -\frac{1}{2}y_L f b_- & -\frac{1}{2}y_L f s_{2h} & \frac{1}{4}y_L f b_+ \\ \frac{\sqrt{5}}{4}y_R f s_{2h} & -M_4 & 0 & 0 & 0 & 0 & 0 \\ -\frac{\sqrt{5}}{4}y_R f s_{2h} & 0 & -M_4 & 0 & 0 & 0 & 0 \\ y_R f (1 - \frac{5}{4}s_h^2) & 0 & 0 & -M_1 & 0 & 0 & 0 \\ \frac{\sqrt{5}}{4}y_R f s_h^2 & 0 & 0 & 0 & -M_9 & 0 & 0 \\ -\frac{\sqrt{5}}{4}y_R f s_h^2 & 0 & 0 & 0 & 0 & -M_9 & 0 \\ \frac{\sqrt{5}}{4}y_R f s_h^2 & 0 & 0 & 0 & 0 & 0 & -M_9 \end{bmatrix}$$

$f, |M_1|, |M_4|, |M_9|, \text{sign}(M_1), \text{sign}(M_4), y_L$ and y_R

Application to the Minimal Composite Higgs Model (MCHM)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

MCHM₅: $T^{(1)}$ $T^{(2)}$ $T^{(3)}$ B $X_{5/3}$

f , $|M_1|$, $|M_4|$, $\text{sign}(M_1)$, y_L and y_R

Fixed by the mass of the top

“Low” scale (HL-LHC):

$$|M_1| \in [0.8, 3.0] \text{ TeV},$$

$$f \in [0.8, 2.0] \text{ TeV},$$

$$M_4 \in [1.2, 3.0] \text{ TeV}$$

$$y_L \in [0.5, 3.0].$$

Region I: $M_1, M_4 > 0$

Region II: $M_1 < 0, M_4 > 0$

“High” scale (FCC and other “future” colliders):

$$|M_1| \in [2, 30] \text{ TeV},$$

$$f \in [0.8, 8.0] \text{ TeV},$$




$$M_4 \in [2, 30] \text{ TeV},$$

$$y_L \in [0.5, 3.0],$$







Clustering for the MCHM₅ (low scale)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

Ideal clustering found for: $N = 11$; variables: $M[t,h]$ & $\theta[t]$



● Region I: $M_1, M_4 > 0$
● Region II: $M_1 < 0, M_4 > 0$




348 parameter space points
 $pp \rightarrow t\bar{t} h h$
(simulated at MG5 @LO)

Clustering for the MCHM₅ (low scale)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

Ideal clustering found for: $N = 11$; variables: $M[t,h]$ & $\theta[t]$












Roughly separates regions
I and II










Clustering for the MCHM₅ (low scale)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

Ideal clustering found for: $N = 11$; variables: $M[t,h]$ & $\theta[t]$



● Region I: $M_1, M_4 > 0$
● Region II: $M_1 < 0, M_4 > 0$



Region I resonances are wider!


Clustering for the MCHM₅ (low scale)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

Ideal clustering found for: $N = 11$; variables: $M[t,h]$ & $\theta[t]$

Usual
simplifying
assumption

Multiple
peaks



Many points have
overlapping top partners!

Clustering for the MCHM₅ (low scale)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

Benchmark points

● Region I: $M_1, M_4 > 0$
● Region II: $M_1 < 0, M_4 > 0$

		C ₁	C ₂	C ₃	C ₄	C ₅	C ₆	C ₇	C ₈	C ₉	C ₁₀	C ₁₁
parameters	M ₁ (GeV)	-1323	-1809	-1483	2965	2882	2999	3000	-1400	-1618	-2384	-2892
	M ₄ (GeV)	1357	1479	2235	1370	1339	1479	1295	1339	1309	1519	1437
	f(GeV)	1199	1593	1071	1393	1220	1168	1484	1265	1229	1110	1646
	y _L	0.91	2.25	1.38	2.35	1.83	2.33	1.98	1.34	1.22	0.51	1.03
	y _R	0.88	0.58	0.72	3.38	3.57	3.28	3.25	0.66	0.74	2.30	0.85
μ(tth) (All Energies)		0.90	0.94	0.86	0.83	0.78	0.79	0.84	0.91	0.90	0.81	0.94
μ(t̄t̄hh) (14 TeV)		2.14	1.47	0.80	1.51	1.53	1.02	2.00	2.25	2.41	1.39	1.58
μ(t̄t̄hh) (100 TeV)		14.58	8.84	3.28	10.28	11.18	7.04	13.42	15.20	16.11	13.68	10.57
NR-t̄t̄hh/t̄t̄hh (14 TeV)		0.37	0.59	0.88	0.45	0.40	0.61	0.35	0.36	0.33	0.46	0.55
NR-t̄t̄hh/t̄t̄hh (100 TeV)		0.05	0.10	0.22	0.07	0.05	0.09	0.05	0.05	0.05	0.05	0.08
M _{T(1)} (TeV)		1.36	1.48	1.66	1.40	1.38	1.51	1.32	1.34	1.31	1.54	1.44
M _{T(2)} (TeV)		1.63	2.02	2.24	3.55	2.61	3.10	3.22	1.61	1.80	1.63	2.20
M _{T(3)} (TeV)		1.79	3.88	2.68	5.55	5.21	4.85	5.67	2.17	2.02	3.47	3.21
M _{B(1)} (TeV)		1.74	3.87	2.68	3.55	2.60	3.10	3.22	2.16	1.99	1.62	2.22
M _{X_{5/3}} (TeV)		1.36	1.48	2.24	1.37	1.34	1.48	1.29	1.34	1.31	1.52	1.44
Γ _{T(1)} (GeV)		8.83	5.49	26.22	51.92	60.01	71.68	44.33	6.44	7.49	43.78	10.63
BR(T ⁽¹⁾ → th)		0.49	0.45	0.31	0.44	0.43	0.42	0.44	0.47	0.47	0.34	0.45
BR(T ⁽¹⁾ → W ⁺ b)		0.018	0	0.47	0.004	0.004	0.003	0.006	0.024	0.016	0.005	0.010
BR(T ⁽¹⁾ → tZ)		0.39	0.41	0.22	0.42	0.43	0.42	0.43	0.40	0.41	0.50	0.41
BR(T ⁽¹⁾ → W ⁺ W ⁻ t)		0.11	0.13	0	0.13	0.13	0.16	0.12	0.10	0.10	0.14	0.12

Clustering for the MCHM₅ (low scale)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

Benchmark points

● Region I: $M_1, M_4 > 0$
● Region II: $M_1 < 0, M_4 > 0$

		C ₁	C ₂	C ₃	C ₄	C ₅	C ₆	C ₇	C ₈	C ₉	C ₁₀	C ₁₁
parameters	M ₁ (GeV)	-1323	-1809	-1483	2965	2882	2999	3000	-1400	-1618	-2384	-2892
	M ₄ (GeV)	1357	1479	2235	1370	1339	1479	1295	1339	1309	1519	1437
	f(GeV)	1199	1593	1071	1393	1220	1168	1484	1265	1229	1110	1646
	y _L	0.91	2.25	1.38	2.35	1.83	2.33	1.98	1.34	1.22	0.51	1.03
	y _R	0.88	0.58	0.72	3.38	3.57	3.28	3.25	0.66	0.74	2.30	0.85
μ(tth) (All Energies)		0.90	0.94	0.86	0.83	0.78	0.79	0.84	0.91	0.90	0.81	0.94
μ(t̄thh) (14 TeV)		2.14	1.47	0.80	1.51	1.55	1.55	1.55	2.41	1.39	1.58	
μ(t̄thh) (100 TeV)		14.58	8.84	3.28	10.28	11.18	7.04	13.42	15.20	16.11	13.68	10.57
NR-t̄thh/t̄thh (14 TeV)		0.37	0.59	0.88	0.45	0.40	0.61	0.35	0.36	0.33	0.46	0.55
NR-t̄thh/t̄thh (100 TeV)		0.05	0.10	0.22	0.07	0.05	0.09	0.05	0.05	0.05	0.05	0.08
M _{T(1)} (TeV)		1.36	1.48	1.66	1.40	1.38	1.51	1.32	1.34	1.31	1.54	1.44
M _{T(2)} (TeV)		1.63	2.02	2.24	3.55	2.61	3.10	3.22	1.61	1.80	1.63	2.20
M _{T(3)} (TeV)		1.79	3.88	2.68	5.55	5.21	4.85	5.67	2.17	2.02	3.47	3.21
M _{B(1)} (TeV)		1.74	3.87	2.68	3.55	2.61	3.10	3.22	1.99	1.99	1.62	2.22
M _{X_{5/3}} (TeV)		1.36	1.48	2.24	1.37	1.31	1.48	1.25	1.31	1.31	1.52	1.44
Γ _{T(1)} (GeV)		8.83	5.49	26.22	51.92	60.01	71.68	44.33	6.44	7.49	43.78	10.63
BR(T ⁽¹⁾ → th)		0.49	0.45	0.31	0.44	0.43	0.42	0.44	0.47	0.47	0.34	0.45
BR(T ⁽¹⁾ → W ⁺ b)		0.018	0	0.47	0.004	0.004	0.003	0.006	0.024	0.016	0.005	0.010
BR(T ⁽¹⁾ → tZ)		0.39	0.41	0.22	0.42	0.43	0.42	0.43	0.40	0.41	0.50	0.41
BR(T ⁽¹⁾ → W ⁺ W ⁻ t)		0.11	0.13	0	0.13	0.13	0.16	0.12	0.10	0.10	0.14	0.12

Stronger mixing

Wider top partner

Clustering for the MCHM₅ (low scale)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

Benchmark points

● Region I: $M_1, M_4 > 0$
● Region II: $M_1 < 0, M_4 > 0$

		C ₁	C ₂	C ₃	C ₄	C ₅	C ₆	C ₇	C ₈	C ₉	C ₁₀	C ₁₁
parameters	M ₁ (GeV)	-1323	-1809	-1483	2965	2882	2999	3000	-1400	-1618	-2384	-2892
	M ₄ (GeV)	1357	1479	2235	1370	1339	1479	1295	1339	1309	1519	1437
	f(GeV)	1199	1593	1071	1393	1220	1168	1484	1265	1229	1110	1646
	y _L	0.91	2.25	1.38	2.35	1.83	2.33	1.98	1.34	1.22	0.51	1.03
	y _R	0.88	0.58	0.72	3.38	3.57	3.28	3.25	0.66	0.74	2.30	0.85
μ(tth) (All Energies)		0.90	0.94	0.86	0.83	0.78	0.79	0.84	0.91	0.90	0.81	0.94
μ(t̄thh) (14 TeV)		2.14	1.47	0.80	1.51	1.53	1.02	2.00	2.25	2.41	1.39	1.58
μ(t̄thh) (100 TeV)		14.58	8.84	3.28	10.28	11.18	7.04	13.42	15.20	16.11	13.68	10.57
NR-t̄thh/t̄thh (14 TeV)		0.37	0.59	0.88	0.45	0.40	0.61	0.35	0.36	0.33	0.46	0.55
NR-t̄thh/t̄thh (100 TeV)		0.05	0.10	0.22	0.07	0.05	0.09	0.05	0.05	0.05	0.05	0.08
M _{T(1)} (TeV)		1.36	1.48	1.66	1.40	1.38	1.51	1.32	1.34	1.31	1.54	1.44
M _{T(2)} (TeV)		1.63	2.02	2.24	3.55	2.61	3.10	3.22	1.61	1.80	1.63	2.20
M _{T(3)} (TeV)		1.79	3.88	2.68	5.55	5.21	4.85	5.67	2.17	2.02	3.47	3.21
M _{B(1)} (TeV)		1.74	3.87	2.68	3.55	2.60	3.10	3.22	2.16	1.99	1.62	2.22
M _{X_{5/3}} (TeV)		1.36	1.48	2.24	1.37	1.34	1.48	1.29	1.34	1.31	1.52	1.44
Γ _{T(1)} (GeV)		8.83	5.49	26.22	51.92	60.01	71.68	44.33	6.44	7.49	43.78	10.63
BR(T ⁽¹⁾ → th)		0.49	0.45	0.31	0.004	0.004	0.003	0.006	0.024	0.016	0.005	0.010
BR(T ⁽¹⁾ → W ⁺ b)		0.018	0	0.47	0.004	0.004	0.003	0.006	0.024	0.016	0.005	0.010
BR(T ⁽¹⁾ → tZ)		0.39	0.41	0.22	0.42	0.43	0.42	0.43	0.40	0.41	0.50	0.41
BR(T ⁽¹⁾ → W ⁺ W ⁻ t)		0.11	0.13	0	0.13	0.13	0.16	0.12	0.10	0.10	0.14	0.12

3-body decays

Only C3 satisfies the usual assumption

Clustering for the MCHM₅ (low scale)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

Benchmark points

● Region I: $M_1, M_4 > 0$
● Region II: $M_1 < 0, M_4 > 0$

		C ₁	C ₂	C ₃	C ₄	C ₅	C ₆	C ₇	C ₈	C ₉	C ₁₀	C ₁₁
parameters	M ₁ (GeV)	-1323	-1809	-1483	2965	2882	2999	3000	-1400	-1618	-2384	-2892
	M ₄ (GeV)	1357	1479	2235	1370	1339	1479	1295	1339	1309	1519	1437
	f(GeV)	1199	1593	1071	1393	1220	1168	1484	1265	1229	1110	1646
	y _L	0.91	2.25	1.38	2.35	1.83	2.33	1.98	1.34	1.22	0.51	1.03
	y _R	0.88	0.58	0.72	3.38	3.57	3.28	3.25	0.66	0.74	2.30	0.85
μ(tth) (All Energies)		0.90	0.94	0.86	0.83	0.78	0.79	0.84	0.91	0.90	0.81	0.94
μ(t̄thh) (14 TeV)		2.14	1.47	0.80	1.51	1.53	1.02	2.00	2.25	2.41	1.39	1.58
μ(t̄thh) (100 TeV)		14.58	8.84	3.28	10.28	11.18	7.04	13.42	15.20	16.11	13.68	10.57
NR-t̄thh/t̄thh (14 TeV)		0.37	0.59	0.88	0.47	0.40	0.61	0.35	0.36	0.33	0.46	0.55
NR-t̄thh/t̄thh (100 TeV)		0.55	0.10	0.22	0.94	0.90	0.69	0.90	0.89	0.89	0.89	0.88
3-body decays The b channel is also mostly suppressed												
M _{T(1)} (TeV)		1.36	1.48	1.66	1.40	1.38	1.51	1.32	1.34	1.31	1.54	1.44
M _{T(2)} (TeV)		1.63	2.02	2.24	3.55	2.61	3.10	3.22	1.61	1.80	1.63	2.20
M _{T(3)} (TeV)		1.79	3.88	2.68	5.55	5.21	4.85	5.67	2.17	2.02	3.47	3.21
M _{B(1)} (TeV)		1.74	3.87	2.68	3.55	2.60	3.10	3.22	2.16	1.99	1.62	2.22
M _{X_{5/3}} (TeV)		1.36	1.48	2.24	1.37	1.34	1.48	1.29	1.34	1.31	1.52	1.44
Γ _{T(1)} (GeV)		8.83	5.49	26.22	51.92	60.01	71.68	44.33	6.44	7.49	43.78	10.63
BR(T ⁽¹⁾ → th)		0.49	0.45	0.31	0.44	0.43	0.42	0.44	0.47	0.47	0.34	0.45
BR(T ⁽¹⁾ → W ⁺ b)		0.018	0	0.47	0.004	0.004	0.003	0.006	0.024	0.016	0.005	0.010
BR(T ⁽¹⁾ → tZ)		0.39	0.41	0.22	0.42	0.43	0.42	0.43	0.40	0.41	0.50	0.41
BR(T ⁽¹⁾ → W ⁺ W ⁻ t)		0.11	0.13	0	0.13	0.13	0.16	0.12	0.10	0.10	0.14	0.12

Clustering for the MCHM₅ (high scale)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

Benchmark points

● Region I: $M_1, M_4 > 0$
● Region II: $M_1 < 0, M_4 > 0$

		E ₁	E ₂	E ₃	E ₄	E ₅	E ₆	E ₇	E ₈	E ₉	E ₁₀
parameters	M ₁ (TeV)	22.7	19.2	11.1	23.0	26.5	3.6	19.3	10.5	-10.7	-27.5
	M ₄ (TeV)	2.4	2.1	3.2	3.2	4.0	22.5	5.1	5.1	25.6	11.3
	f (GeV)	1913	3273	7144	1190	1300	1711	1288	2812	2432	1412
	y _L	2.45	0.87	2.85	2.43	0.99	2.00	2.35	1.84	2.57	1.73
	y _R	1.10	1.24	2.01	1.54	3.53	1.31	2.35	3.13	1.11	2.96
μ(<i>tth</i>) (All Energies)		0.95	0.97	0.99	0.88	0.83	0.94	0.88	0.97	0.96	0.90
μ(<i>tthh</i>) (100 TeV)		1.26	1.91	1.03	0.82	0.81	0.86	0.75	0.91	0.92	0.78
NR- <i>tthh</i> / <i>tthh</i> (100 TeV)		0.71	0.48	0.95	0.90	0.82	1.00	1.00	1.02	1.01	1.01
M _{T(1)} (TeV)		2.45	2.12	3.21	3.23	4.07	4.28	5.08	5.15	11.0	11.3
M _{T(2)} (TeV)		5.27	3.55	18.1	4.32	4.28	22.5	5.90	7.31	25.6	11.6
M _{T(3)} (TeV)		22.8	19.7	20.6	23.1	26.9	22.8	19.5	13.7	26.4	27.8
M _{B(1)} (TeV)		5.28	3.55	20.6	4.33	4.24	22.8	5.90	7.30	26.4	11.6
M _{X_{5/3}} (TeV)		2.44	2.11	3.20	3.22	4.04	22.5	5.06	5.14	25.6	11.3
Γ _{T(1)} (TeV)		0.04	0.04	0.08	0.14	0.96	0.28	0.76	0.84	1.22	8.97
Γ _{T(1)} /M _{T(1)}		1.6%	1.9%	2.5%	4.3%	24%	6.5%	15%	16%	11%	79%
BR(T ⁽¹⁾ → th)		0.35	0.38	0.29	0.29	0.15	0.26	0.18	0.17	0.25	0.05
BR(T ⁽¹⁾ → W ⁺ b)		0.003	0.004	0	0.001	0	0.50	0	0	0.50	0
BR(T ⁽¹⁾ → tZ)		0.34	0.37	0.28	0.28	0.33	0.25	0.18	0.18	0.25	0.06
BR(T ⁽¹⁾ → W ⁺ W ⁻ t)		0.30	0.25	0.43	0.43	0.52	0	0.64	0.65	0	0.89

Clustering for the MCHM₅ (high scale)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

Benchmark points

● Region I: $M_1, M_4 > 0$
● Region II: $M_1 < 0, M_4 > 0$

		E ₁	E ₂	E ₃	E ₄	E ₅	E ₆	E ₇	E ₈	E ₉	E ₁₀
Parameters	M ₁ (TeV)	22.7	19.2	11.1	23.0	26.5	3.6	19.3	10.5	-10.7	-27.5
	M ₄ (TeV)	2.4	2.1	3.2	3.2	4.0	22.5	5.1	5.1	25.6	11.3
	f (GeV)	1913	3273	7144	1190	1300	1711	1288	2812	2432	1412
	3-body decays										
	γ_R	2.45	0.87	2.85	2.43	0.99	2.00	2.35	1.84	2.57	1.73
	$\mu(t\bar{t}h)$ (All Energies)	1.10	1.24	2.01	1.54	3.53	1.31	2.35	3.13	1.11	2.96
	$\mu(t\bar{t}hh)$ (100 TeV)	0.95	0.97	0.99	0.88	0.83	0.94	0.88	0.97	0.96	0.90
	NR- $t\bar{t}hh/t\bar{t}hh$ (100 TeV)	1.26	1.91	1.03	0.82	0.81	0.86	0.75	0.91	0.92	0.78
	$M_{T(1)}$ (TeV)	0.71	0.48	0.95	0.90	0.82	1.00	1.00	1.02	1.01	1.01
	$M_{T(2)}$ (TeV)	2.45	2.12	3.21	3.23	4.07	4.28	5.08	5.15	11.0	11.3
	$M_{T(3)}$ (TeV)	5.27	3.55	18.1	4.32	4.28	22.5	5.90	7.31	25.6	11.6
	$M_{B(1)}$ (TeV)	22.8	19.7	20.6	23.1	26.9	22.8	19.5	13.7	26.4	27.8
	$M_{X_{5/3}}$ (TeV)	5.28	3.55	20.6	4.33	4.24	22.8	5.90	7.30	26.4	11.6
	$\Gamma_{T(1)}$ (TeV)	2.44	2.11	3.20	3.22	4.04	22.5	5.06	5.14	25.6	11.3
	$\Gamma_{T(1)}/M_{T(1)}$	0.04	0.04	0.08	0.14	0.96	0.28	0.76	0.84	1.22	8.97
	BR(T ⁽¹⁾ → th)	1.6%	1.9%	2.5%	4.3%	24%	6.5%	15%	16%	11%	79%
	BR(T ⁽¹⁾ → W ⁺ b)	0.35	0.38	0.29	0.29	0.15	0.26	0.18	0.17	0.25	0.05
	BR(T ⁽¹⁾ → tZ)	0.003	0.004	0	0.001	0	0.50	0	0	0.50	0
	BR(T ⁽¹⁾ → W ⁺ W ⁻ t)	0.34	0.37	0.28	0.28	0.33	0.25	0.18	0.18	0.25	0.06
		0.30	0.25	0.43	0.43	0.52	0	0.64	0.65	0	0.89

Clustering for the MCHM₅ (high scale)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

Benchmark points

● Region I: $M_1, M_4 > 0$
● Region II: $M_1 < 0, M_4 > 0$

		E ₁	E ₂	E ₃	E ₄	E ₅	E ₆	E ₇	E ₈	E ₉	E ₁₀
Parameters	M ₁ (TeV)	22.7	19.2	11.1	23.0	26.5	3.6	19.3	10.5	-10.7	-27.5
	M ₄ (TeV)	2.4	2.1	3.2	3.2	4.0	22.5	5.1	5.1	25.6	11.3
	f (GeV)	1913	3273	7144	1190	1300	1711	1288	2812	2432	1412
	γ_R	2.45	0.87	2.85	2.43	0.99	2.00	2.35	1.84	2.57	1.73
3-body decays		1.10	1.24	2.01	1.54	3.53	1.31	2.35	3.13	1.11	2.96
$\mu(t\bar{t}h)$ (All Energies)		0.95	0.97	0.99	0.88	0.83	0.94	0.88	0.97	0.96	0.90
$\mu(t\bar{t}hh)$ (100 TeV)		1.26	1.91	1.03	0.82	0.81	0.86	0.75	0.91	0.92	0.78
NR- $t\bar{t}hh/t\bar{t}hh$ (100 TeV)		0.71	0.48	0.95	0.90	0.82	1.00	1.00	1.02	1.01	1.01
$M_{T(1)}$ (TeV)		2.45	2.12	3.21	3.23	4.07	4.28	5.08	5.15	11.0	11.3
$M_{T(2)}$ (TeV)		5.27	3.55	18.1	4.32	4.28	22.5	5.90	7.31	25.6	11.6
$M_{T(3)}$ (TeV)		22.8	19.7	20.6	23.1	26.9	22.8	19.5	13.7	26.4	27.8
$M_{B(1)}$ (TeV)		5.28	3.55	20.6	4.33	4.24	22.8	5.90	7.30	26.4	11.6
$M_{X_{5/3}}$ (TeV)		2.44	2.11	3.20	3.22	4.04	22.5	5.06	5.14	25.6	11.3
$\Gamma_{T(1)}$ (TeV)		0.04	0.04	0.08	0.14	0.90	0.28	0.76	0.84	1.22	8.97
$\Gamma_{T(1)}/M_{T(1)}$		1.6%	1.9%	2.5%	4.3%	24%	6.5%	15%	16%	11%	79%
BR(T ⁽¹⁾ → th)		0.35	0.38	0.29	0.29	0.13	0.26	0.18	0.17	0.25	0.05
BR(T ⁽¹⁾ → W ⁺ b)		0.003	0.004	0	0.001	0	0.50	0	0	0.50	0
BR(T ⁽¹⁾ → tZ)		0.34	0.37	0.28	0.28	0.33	0.25	0.18	0.18	0.25	0.06
BR(T ⁽¹⁾ → W ⁺ W ⁻ t)		0.30	0.25	0.43	0.43	0.52	0	0.64	0.65	0	0.89

Clustering for the MCHM₅ (high scale)

C. Bautista, L. de Lima, R. D. Matheus, E. Pontón, L. A. F. do Prado, A Savoy-Navarro. JHEP 2021, 49 (2021)

Benchmark points

● Region I: $M_1, M_4 > 0$
● Region II: $M_1 < 0, M_4 > 0$

		E ₁	E ₂	E ₃	E ₄	E ₅	E ₆	E ₇	E ₈	E ₉	E ₁₀
Parameters	M_1 (TeV)	22.7	19.2	11.1	23.0	26.5	3.6	19.3	10.5	-10.7	-27.5
	M_4 (TeV)	2.4	2.1	3.2	3.2	4.0	22.5	5.1	5.1	25.6	11.3
	f (GeV)	1913	3273	7144	1190	1300	1711	1288	2812	2432	1412
	γ_R	2.45	0.87	2.85	2.43	0.99	2.00	2.35	1.84	2.57	1.73
	γ_R	1.10	1.24	2.01	1.54	3.53	1.31	2.35	3.13	1.11	2.96
$\mu(t\bar{t}h)$ (All Energies)		0.95	0.97	0.99	0.88	0.83	0.94	0.88	0.97	0.96	0.90
$\mu(t\bar{t}hh)$ (100 TeV)		1.26	1.91	1.03	0.82	0.81	0.86	0.75	0.91	0.92	0.78
NR- $t\bar{t}hh/t\bar{t}hh$ (100 TeV)		0.71	0.48	0.95	0.90	0.82	1.00	1.00	1.02	1.01	1.01
$M_{T(1)}$ (TeV)		2.45	2.12	3.21	3.23	4.07	4.28	5.08	5.15	11.0	11.3
$M_{T(2)}$ (TeV)		5.27	3.55	18.1	4.32	4.28	22.5	5.90	7.31	25.6	11.6
$M_{T(3)}$ (TeV)		22.8	19.7	20.6	23.1	26.9	22.8	19.5	13.7	26.4	27.8
$M_{B(1)}$ (TeV)		5.28	3.55	20.6	4.33	4.24	22.8	5.90	7.30	26.4	11.6
$M_{X_{5/3}}$ (TeV)		2.44	2.11	3.20	3.22	4.04	22.5	5.06	5.14	25.6	11.3
$\Gamma_{T(1)}$ (TeV)		0.04	0.04	0.08	0.14	0.96	0.28	0.76	0.84	1.22	8.97
$\Gamma_{T(1)}/M_{T(1)}$		1.6%	1.9%	2.5%	4.3%	24%	6.5%	15%	16%	11%	79%
BR($T^{(1)} \rightarrow th$)		0.35	0.38	0.29	0.29	0.15	0.26	0.18	0.17	0.25	0.05
BR($T^{(1)} \rightarrow W^+b$)		0.003	0.004	0	0.001	0	0.50	0	0	0.50	0
BR($T^{(1)} \rightarrow tZ$)		0.34	0.37	0.28	0.28	0.33	0.25	0.18	0.18	0.25	0.06
BR($T^{(1)} \rightarrow W^+W^-t$)		0.30	0.25	0.43	0.43	0.52	0	0.64	0.65	0	0.89


Conclusions

- We have to look far from the street lamp now
- “Smart” algorithms can help to **understand** complete models
- Regarding top partners:
 - Three body decays are important
 - A simplified hierarchical spectrum is only realized in small regions of the MCHM



Extras

Wider resonances in Region I



Must interfere negatively as mass must vanish in the SO(5) restoration limit ($M_1 - M_4$)

$$m_t \sim y_L y_R |M_4 - M_1|$$

In region I, both parameters are positive, and this can be small

To compensate, and obtain the correct m_t , the couplings must be larger, leading to wider resonances

Higgs Sector

$$U = \begin{pmatrix} \mathbb{1}_{3 \times 3} & \vec{0} & \vec{0} \\ \vec{0}^T & \cos \frac{h_0+h}{f} & \sin \frac{h_0+h}{f} \\ \vec{0}^T & -\sin \frac{h_0+h}{f} & \cos \frac{h_0+h}{f} \end{pmatrix},$$

$$f \sin \frac{h_0}{f} \equiv v = 246 \text{ GeV}$$

$$\xi = \frac{v^2}{f^2} = \sin^2 \frac{h_0}{f}$$

MCHM₅

$$\mathcal{L} = \mathcal{L}_{\text{elem}} + \mathcal{L}_{\text{comp}}^5 + \mathcal{L}_{\text{mix}}^5 + \mathcal{L}_{\text{int}}^5$$

$$\mathcal{L}_{\text{elem}} = \bar{q}_L i \not{D} q_L + \bar{t}_R i \not{D} t_R$$

$$\Psi_1 \sim \tilde{T}$$

$$\Psi_4 \sim (X_{5/3}, X_{2/3}, T, B)$$

$$\mathcal{L}_{\text{comp}}^5 = \bar{\Psi}_4 i (\not{D} - i\phi) \Psi_4 - M_4 \bar{\Psi}_4 \Psi_4 + \bar{\Psi}_1 i \not{D} \Psi_1 - M_1 \bar{\Psi}_1 \Psi_1$$

$$\begin{aligned} \mathcal{L}_{\text{mix}}^5 &= f \bar{Q}_L^5 U [y_{L4} \Psi_4 + y_{L1} \Psi_1] + \text{h.c.} \\ &+ f \bar{T}_R^5 U [y_{R4} \Psi_4 + y_{R1} \Psi_1] + \text{h.c.} \end{aligned}$$

$$\mathcal{L}_{\text{int}}^5 = -i c_L \bar{\Psi}_4 P_L \not{d} \Psi_1 - i c_R \bar{\Psi}_4 P_R \not{d} \Psi_1 + \text{h.c.}$$

MCHM₁₄

$$\mathcal{L} = \mathcal{L}_{\text{elem}} + \mathcal{L}_{\text{comp}}^{14} + \mathcal{L}_{\text{mix}}^{14} + \mathcal{L}_{\text{int}}^{14}$$

$$\Psi_9 \sim (U_{8/3}, U_{5/3}, U_{2/3}, V_{5/3}, V_{2/3}, V_{-1/3}, F_{2/3}, F_{-1/3}, F_{-4/3})$$



$$\mathcal{L}_{\text{comp}}^{14} = \mathcal{L}_{\text{comp}}^5 + \text{Tr} \left[\bar{\Psi}_9 (i \not{D} \Psi_9 - i [\not{e}, \Psi_9]) \right] - M_9 \text{Tr} \left[\bar{\Psi}_9 \Psi_9 \right]$$

$$\begin{aligned} \mathcal{L}_{\text{mix}}^{14} = & f \text{Tr} \left[U^\top \bar{Q}_L^{14} U (y_{L9} \Psi_9 + y_{L4} \Psi_4 + y_{L1} \Psi_1) \right] + \text{h.c.} \\ & + f \text{Tr} \left[U^\top \bar{T}_R^{14} U (y_{R9} \Psi_9 + y_{R4} \Psi_4 + y_{R1} \Psi_1) \right] + \text{h.c.} \end{aligned}$$

$$\mathcal{L}_{\text{int}}^{14} = -i c_4 \bar{\Psi}_4 \not{D} \Psi_1 - i c_9 \bar{\Psi}_9^{ij} \not{D}^i \Psi_4^j - i \frac{c_{T9}}{4\pi f} \bar{\Psi}_9^{ij} d_\mu^i d^{j\mu} \tilde{T} + \text{h.c.}$$

Constraints MCHM₅

Overlaid regions indicate constraints: the dark one is given by direct exclusion of top partners in the left plot and by expected constraints in the HL-LHC in the right one ($M_{T^{(1)}} < 4$ TeV). In the white region, the top mass cannot be reached without violating perturbativity.





$$0.69 \leq c_g \leq 1.33$$




$$0.33 \leq \mu(t\bar{t}h) \leq 2.07$$

Constraints MCHM₁₄

Overlaid regions indicate constraints: the dark one is given by direct exclusion of top partners in the left plot and by expected constraints in the HL-LHC in the right one ($M_{T(1)} < 4$ TeV). In the white region, the top mass cannot be reached without violating perturbativity. The green region is constrained by c_g measurements in the left and by the c_g expected constraints in the right.









tthh process



tth

$M_{\Upsilon^{(1)}} [\text{TeV}]$



NR-tthh

	P ₁	P ₂	P ₃	P ₄	P ₅	Disregarded diagrams
$\sigma_{\mathcal{W}h}/\sigma_{\text{NR}}^{\bar{t}t\bar{t}h}$ (14 TeV) $\sigma_{\mathcal{W}h}/\sigma_{\text{NR}}^{\bar{t}t\bar{t}h}$ (100 TeV)	1.05 1.05	1.04 1.03	1.03 1.03	1.01 1.01	1.01 1.01	
$\sigma_{\text{Yuk}}/\sigma_{\text{NR}}^{\bar{t}t\bar{t}h}$ (14 TeV) $\sigma_{\text{Yuk}}/\sigma_{\text{NR}}^{\bar{t}t\bar{t}h}$ (100 TeV)	0.86 0.87	0.85 0.87	0.84 0.87	0.82 0.85	0.82 0.85	
$\sigma_{\text{NR}}^{\bar{t}t\bar{t}h}/\sigma_{\text{SM}}^{\bar{t}t\bar{t}h}$ (14 TeV) $\sigma_{\text{NR}}^{\bar{t}t\bar{t}h}/\sigma_{\text{SM}}^{\bar{t}t\bar{t}h}$ (100 TeV) $(y_t/y_t^{\text{SM}})^4$	0.65 0.65 0.69	0.69 0.69 0.72	0.82 0.82 0.85	0.94 0.93 0.95	0.90 0.89 0.91	

Table 3. Study of NR- $\bar{t}t\bar{t}h$ for the MCHM₅ points in table 1. The cross sections $\sigma_{\mathcal{W}h}$ and σ_{Yuk} are obtained by disregarding the classes of diagrams on the last column and σ_{NR} is the total NR- $\bar{t}t\bar{t}h$. The LO SM $\bar{t}t\bar{t}h$ production is indicated by σ^{SM} and $\sigma_{\text{Yuk}}^{\text{SM}}$ means we disregarded the SM trilinear Higgs coupling. The top Yukawa couplings are indicated by y_t and y_t^{SM} in the MCHM and SM respectively.

EFT

$$\mathcal{L}_h = \frac{1}{2} \partial_\mu h \partial^\mu h - \frac{1}{2} m_h^2 h^2 - \kappa_\lambda \lambda_{\text{SM}} v h^3 - \frac{m_t}{v} \left(v + \kappa_t h + \frac{c_2}{v} h h \right) (\bar{t}_L t_R + \text{h.c.})$$

$$+ \frac{1}{4} \frac{\alpha_s}{3\pi v} \left(c_g h - \frac{c_{2g}}{2v} h h \right) G^{\mu\nu} G_{\mu\nu}$$




Figure 20. Values of some selected EFT parameters in the low scale scan of the MCHM₅ and the MCHM₁₄ parameter spaces. The colors indicate the different Regions in each model (I and II for the MCHM₅ and I, II, III and IV for the MCHM₁₄, in that order). The SM is represented by the black square.